Summary of Water Conditions

May 1, 2018

April precipitation in Northern California was above average, most of which was generated by a strong atmospheric river event during the first part of the month which prematurely melted some of the snowpack and forced some foothill reservoir operators into the flood control mode of operation. The water supply outlook improved moderately during the month.

Forecasts of median April through July runoff went up about 10 percent from April 1 and are now expected to be about 80 percent of average, and 75 percent for the water year. Runoff from snowmelt last year was about 190 percent of average.

Snowpack water content shrank a lot during April due to a relatively warm major storm, and is now about 25 percent of the May 1 average. Most lower elevation courses are bare.

Precipitation from October through April was 75 percent of average compared to 70 percent one month ago and 170 percent last year. The range is from about 90 percent in the north to 30-40 percent in the south.

Runoff to date has been about 75 percent of average compared to 235 percent last year at this time. Runoff in April was estimated at 135 percent, including some early snowmelt. Estimated runoff of the eight major rivers of the Sacramento-San Joaquin river region in April was 4.24 million acre-feet in April.


Reservoir storage which includes carryover storage from 2017, is good at about 110 percent of average and is about the same as one year ago.

SUMMARY OF WATER CONDITIONS IN PERCENT OF AVERAGE

HYDROLOGIC REGION	PRECIPITATION OCTOBER 1 TO DATE	MAY 1 SNOW WATER CONTENT	MAY 1 RESERVOIR STORAGE	RUNOFF OCTOBER 1 TO DATE	APRIL-JULY RUNOFF FORECAST	WATER YEAR RUNOFF FORECAST
NORTH COAST	80	10	95	55	45	40
SAN FRANCISCO BAY	70		90	30		
CENTRAL COAST	60		60	40		
SOUTH COAST	40		80	15		
SACRAMENTO RIVER	90	25	105	75	80	70
SAN JOAQUIN RIVER	85	40	130	110	85	85
TULARE LAKE	70	30	145	90	70	70
NORTH LAHONTAN	90	30	170	145	100	105
SOUTH LAHONTAN	50		115	105	75	85
COLORADO RIVER	30					
STATEWIDE	75	25	110	75	80	75


CALIFORNIA COOPERATIVE SNOW SURVEYS SEASONAL PRECIPITATION

IN PERCENT OF AVERAGE TO DATE October 1, 2017 through April 30, 2018



CR - Colorado River-Desert

DEPARMENT OF WATER RESOURCES CALIFORNIA COOPERATIVE SNOW SURVEYS FORECAST OF APRIL-JULY UNIMPAIRED SNOWMELT RUNOFF May 1, 2018



^{*} FORECAST BY DEPARTMENT OF WATER AND POWER, CITY OF LOS ANGELES

May 1, 2018 FORECASTS **APRIL-JULY UNIMPAIRED RUNOFF**

HYDROLOGIC REGION	н	ا Apr-Jul I ISTORICAL		Runoff in 1,000 Acre-Feet (1) FORECAST			
and Watershed	50 Yr	Max of	Min of	Apr-Jul	Pct	80%	
	Avg (2)	Record (10)	Record (10)	Forecast	of Avg	Probability Range (1)	
North Coast	(2)	(10)	(10)	<u>;</u>	Avy :	range (1)	
Trinity River at Lewiston Lake	639	1,593	80	300	47%	250 - 350	
SACRAMENTO RIVER		_,					
Upper Sacramento River							
Sacramento River at Delta above Shasta Lake	295	751	39	175	59%		
McCloud River above Shasta Lake	385	850	185	320	83%		
Pit River near Montgomery Creek + Squaw Creek	1,020	2,098	480	890	87%		
Total Inflow to Shasta Lake	1,756	3,525	711	1,440	82%	1,220 - 1,610	
Sacramento River above Bend Bridge, near Red Bluff	2,421	5,117	943	1,940	80%	1,620 - 2,260	
Feather River							
Feather River at Lake Almanor near Prattville (3)	333	675	120	250	75%		
North Fork at Pulga (3)	1,028	2,416	243	740	72%		
Middle Fork near Clio (4) South Fork at Ponderosa Dam (3)	86 110	518 267	4 13	59 74	69% 67%		
Feather River at Oroville	1,704	4,676	378	1,310	77%	1,080 - 1,480	
Yuba River	1,704	4,070	370	1,310	1170	1,000 1,400	
North Yuba below Goodyears Bar	279	647	51	230	82%		
Inflow to Jackson Mdws and Bowman Reservoirs (3)	112	236	25	90	80%		
South Yuba at Langs Crossing (3)	233	481	57	180	77%		
Yuba River near Smartsville plus Deer Creek	968	2,424	151	810	84%	670 - 950	
American River							
North Fork at North Fork Dam (3)	262	716	43	0	0%		
Middle Fork near Auburn (3)	522	1,406	100	460	88%		
Silver Creek below Camino Diversion Dam (3)	173	386	37	150	87%		
American River below Folsom Lake	1,199	3,074	185	1,100	92%	910 - 1,300	
SAN JOAQUIN RIVER							
Cosumnes River at Michigan Bar	125	446	8	130	104%	100 - 165	
Mokelumne River							
North Fork near West Point (5)	437	829	104	0	0%		
Total Inflow to Pardee Reservoir	457	1,076	75	390	85%	350 - 450	
Stanislaus River	004	700	6.4	070	040/		
Middle Fork below Beardsley Dam (3)	334	702	64	270	81%		
North Fork Inflow to McKays Point Dam (3) Stanislaus River below Goodwin Reservoir (9)	224 682	503 1,710	34 116	185 580	83% 85%	510 - 680	
Tuolumne River	002	1,710	110	300	0370	310 000	
Cherry Creek & Eleanor Creek near Hetch Hetchy	315	727	97	280	89%		
Tuolumne River near Hetch Hetchy	604	1,392	153	540	89%		
Tuolumne River below La Grange Reservoir (9)	1,193	2,682	301	1,050	88%	930 - 1,220	
Merced River				,			
Merced River at Pohono Bridge	372	888	80	310	83%		
Merced River below Merced Falls (9)	623	1,588	104	500	80%	440 - 570	
San Joaquin River							
San Joaquin River at Mammoth Pool (7)	1,026	2,279	235	850	83%		
Big Creek below Huntington Lake (8)	91	264	11	72	79%		
South Fork near Florence Lake (7)	201	511	58	175	87%		
San Joaquin River inflow to Millerton Lake	1,228	3,355	193	1,000	81%	860 - 1,140	
TULARE LAKE							
Kings River							
North Fork Kings River near Cliff Camp (3)	239	565	50	190	79%		
Kings River below Pine Flat Reservoir	1,210	3,113	208	960	79%	820 - 1,100	
Kaweah River below Terminus Reservoir	285	814	42	165	58%	140 - 190	
Tule River below Lake Success	63	259	1	28	44%	22 - 35	
Kern River							
Kern River near Kernville	384	1,203	83	220	57%	046	
Kern River inflow to Lake Isabella	458	1,657	57	250	55%	210 - 300	

⁽¹⁾ See inside the back cover for definition.
(2) All 50 year averages are based on years 1966-2015 unless otherwise noted.

^{(3) 50} year average based on years 1941-90. (4) 44 year average based on years 1936-79.

^{(5) 36} year average based on years 1936-72.(6) 45 year average based on years 1936-81.(7) 50 year average based on years 1953-2002.(8) 50 year average based on years 1946-1995.

May 1, 2018 FORECASTS WATER YEAR UNIMPAIRED RUNOFF

н	ISTORICAI			V	Vater Ye		paired F	Runoff in	1,000 A	cre-Fee	t (1)	FORECAST			
50 Yr Avg (2)	Max of Record (10)	Min of Record (10)	Oct Thru Jan	Feb *	Mar *	Apr *	May	Jun	Jul	Aug	Sep	Water Year Forecast	Pct of Avg	80% Probab Range	ility
1,348	2,990	200	135	43	91	156	105	31	8	2	0	571	42%	520 -	625
860 1,183 3,002 5,831 8,544	1,966 2,353 5,150 10,796 17,180	165 557 1,484 2,479 3,294	1,156 1,582	254 325	636 848	533 799	425 525	270 350	212 266	195 231	189 229	3,870 5,155	66% 60%	3,575 - 4,735 -	4,090 5,555
780 2,417 219 291 4,407	1,269 4,400 637 562 10,178	366 666 24 32 995	843	181	867	802	290	135	83	66	59	3,325	75%	3,040 -	3,520
564 181 379 2,268	1,056 292 565 5,604	102 30 98 369	488	88	514	510	225	60	15	8	8	1,915	84%	1,770 -	2,060
616 1,070 318 2,626	1,234 2,575 705 7,391	66 144 59 349	502	98	661	713	285	88	14	3	3	2,367	90%	2,175 -	2,570
379	1,253	20	43	9	142	102	22	5	1	0	0	324	86%	290 -	360
626 748	1,009 1,901	197 129	106	18	127	205	140	40	5	2	1	644	86%	600 -	705
471	929	88	454	00	407	045	100	70	4.5				0.407	000	4 070
1,149 461 770 1,909	3,078 1,147 1,661 4,631	155 123 258 383	154 229	29 34	197 357	315 474	180 345	70 200	15 31	10	2	965 1,685	84%	892 <i>-</i> 1,560 <i>-</i>	1,070 1,860
461 992	1,020 2,787	92 150	73	16	185	249	170	67	14	5	1	780	79%	715 -	860
1,337 112 248 1,793	2,964 298 653 4,642	308 14 71 327	130	27	211	380	360	210	50	18	8	1,395	78%	1,245 -	1,550
284 1,702 451 147	607 4,287 1,402 615	58 359 89 10	116 28 13	23 7 3	164 46 21	362 86 16	345 53 10	205 22 2	48 4 0	19 2 0	11 1 0	1,295 249 64	76% 55% 44%	1,145 - 220 - 55 -	1,440 280 75
558 728	1,577 2,318	163 130	103	18	48	84	80	63	23	12	9	440	60%	395 -	500

⁽⁹⁾ Forecast point names based on USGS gage names. Stanislaus below Goodwin also known as inflow to New Melones, Tuolumne River below La Grange also known as inflow to Don Pedro, Merced River below Merced Falls also known as inflow to McClure.
(10) For the tributaries, the period of record over which the minimum and maximum values are found does not include years after water year 2011.
* Unimpaired runoff in months prior to forecast date are based on measured flows.


May 1, 2018 FORECASTS **APRIL-JULY UNIMPAIRED RUNOFF**

HYDROLOGIC REGION		Jnimpaired F		000 Acre-Feet (1) FORECAST		
and Watershed	50 Yr Avg (2)	Max of Record (6)	Min of Record (6)	Apr-Jul Forecast	Pct of Avg	
NORTH COAST						
Scott River Scott River nr Ft Jones (3)	173	398	22	103	60%	
Klamath River Total inflow to Upper Klamath Lake (4)	475	1,150	149	311	65%	
NORTH LAHONTAN						
Truckee River Lake Tahoe to Farad accretions Lake Tahoe Rise (assuming gates closed, ft)	250 1.3	713 5.4	48 0.2		104% 97%	
Carson River West Fork Carson River at Woodfords East Fork Carson River near Gardnerville	52 182	135 480	10 43		104% 107%	
Walker River West Walker River below Little Walker, near Coleville East Walker River near Bridgeport	153 61	410 209	35 7	140 50	92% 82%	
SOUTH LAHONTAN						
Owens River Total tributary flow to Owens River (5)	231	579	84	170	73%	


⁽¹⁾ See inside the back cover for definition.

⁽²⁾ All 50 year averages are based on years 1966-2015 unless otherwise noted.
(3) Forecast by National Weather Service California-Nevada River Forecast Center. 30 yr average (1981-2010).
(4) Forecast by U.S. Natural Resources Conservation Service and National Weather Service California-Nevada River Forecast Center, April through (4) Forecast by 0.3.1 Natural resolutes conservation service and value and value as service canonical revolution for the conservation service and value as service canonical revolution for the conservation of the conservation service and value as service canonical revolution for the conservation of the conservation service and value as service canonical revolution for the conservation service and value as service canonical revolution for the conservation service and value as service canonical revolution for the conservation service and value as service canonical revolution for the conservation service and value as service canonical revolution for the conservation service and value as service canonical revolution for the conservation service and value as service canonical revolution for the conservation service and value as service canonical revolution for the conservation service and value as service canonical revolution for the conservation for the conservation service and value as service canonical revolution for the conservation for the conservation service and value as service canonical revolution for the conservation for


NORTH COAST REGION




SNOWPACK First of the month measurements made at 5 snow courses indicate an area wide snow water equivalent of 2.2 inches. This is 5 percent of the seasonal April 1 average and 10 percent of the May 1 average165. Last year this time the pack was holding 39.1 inches of water.



PRECIPITATION Seasonal precipitation (October 1 through to the end of April) on this area was 80 percent of normal. Precipitation last month was about 165 percent of the monthly average. Season precipitation at this time last year stood at 160 percent of normal.




RESERVOIR STORAGE First of the month storage at 6 reservoirs was 2.35 million acre-feet which is 95 percent of average. About 75 percent of available capacity was being used. Storage in these reservoirs at this time last year was 115 percent of average.




RUNOFF Seasonal runoff of streams draining this area totaled 5.85 million acre-feet which is 55 percent of average. Last year, runoff for the same period was 200 percent of average.

SACRAMENTO RIVER REGION





SNOWPACK First of the month measurements made at 62 snow courses indicate an area wide snow water equivalent of 6.7 inches. This is 15 percent of the seasonal April 1 average and 25 percent of the May 1 average. Last year this time the pack was holding less than 40.9 inches of water.

PRECIPITATION Seasonal precipitation (October 1 through to the end of April) on this area was 90 percent of normal. Precipitation last month was about 145 percent of the monthly average. Season precipitation at this time last year stood at 190 percent of normal.



Reservoir Storage Contents of major reservoirs in % of capacity






RESERVOIR STORAGE First of the month storage at 43 reservoirs was 13.68 million acre-feet which is 105 percent of average. About 85 percent of available capacity was being used. Storage in these reservoirs at this time last year was 110 percent of average.

RUNOFF Seasonal runoff of streams draining this area totaled 9.82 million acre-feet which is 75 percent of average. Last year, runoff for the same period was 240 percent of average.

The Sacramento Region 40-30-30 Water Supply Index is forecast to be 7.2 assuming median meteorological conditions for the remainder of the year. This classifies the year as "below normal" in the Sacramento Valley according to the State Water Resources Control Board.

SAN JOAQUIN RIVER AND TULARE LAKE REGIONS



Kings

San Joaquin

WY 2017

Kaweah

WY 2018

Merced

Tuolumne

100

SNOWPACK- First of the month measurements made at 55 San Joaquin Region snow courses indicate an area wide snow water equivalent of 12 inches. This is 30 percent of the seasonal April 1 average and 40 percent of the May 1 average. Last year this time the pack was holding 53.6 inches of water. At the same time 32 Tulare Lake snow courses indicate a basin-wide snow water equivalent of less than 6.9 inches. This is 20 percent of the seasonal April 1 average and 30 percent of the May 1 average. Last year this time the pack was holding 41.2 inches of water.

PRECIPITATION- Seasonal precipitation (October 1 through to the end of April) on the **San Joaquin Region** was 85 percent of normal. Precipitation last month was about 120 percent of the monthly average. Season precipitation at this time last year stood at 185 percent of normal. Seasonal precipitation (October 1 through to the end of April) on the **Tulare Lake Region** was 70 percent of normal. Precipitation last month was about 80 percent of the monthly average. Season precipitation at this time last year stood at 175 percent of normal.


RESERVOIR STORAGE- First of the month storage in 34 San Joaquin Region reservoirs was 9.87 million acre-feet which is 130 percent of average. About 85 percent of available capacity was being used. Storage in these reservoirs at this time last year was 120 percent of average. First of the month storage in 6 **Tulare Lake Region** reservoirs was 1.51 million acre-feet which is 145 percent of average. About 70 percent of available capacity was being used. Storage in these reservoirs at this time last year was 100 percent of average.

RUNOFF- Seasonal runoff of streams draining the **San Joaquin Region** totaled 3.81 million acre-feet which is 110 percent of average. Last year, runoff for the same period was 305 percent of average. Seasonal runoff of streams draining the **Tulare Lake Region** area totaled 1.14 million acre-feet which is 90 percent of average. Last year, runoff for the same period was 265 percent of average.

The San Joaquin Region 60-20-20 Water Supply Index is forecast to be 3.0 assuming 75 percent of median meteorological conditions. This classifies the year as "Below Normal" in the San Joaquin according to the State Water Resources Control Board.

Kern

NORTH AND SOUTH LAHONTAN REGIONS




SNOWPACK First of the month measurements made at 4 **North Lahontan Region** snow courses indicate an area wide snow water equivalent of less than 7.4 inches. This is 25 percent of the seasonal April 1 average and 30 percent of the May 1 average. Last year this time the pack was holding 34.4 inches of water. At the same time 6 **South Lahontan Region** snow sensors indicate a basin-wide snow water equivalent of 4.9 inches. This is 20 percent of the seasonal April 1 average and 25 percent of the May 1 average. Last year this time the pack was holding 27.6 inches of water.

PRECIPITATION Seasonal precipitation (October 1 through to the end of April) on the **North Lahontan Region** was 90 percent of normal. Precipitation last month was about 145 percent of the monthly average. Season precipitation at this time last year stood at 210 percent of normal. Seasonal precipitation (October 1 through to the end of April) on the **South Lahontan Region** was 50 percent of normal. Precipitation last month was about 30 percent of the monthly average. Season precipitation at this time last year stood at 140 percent of normal.

RESERVOIR STORAGE First of the month storage in 5 **North Lahontan Region** reservoirs was 1.00 million acre-feet which is 170 percent of average. About 95 percent of available capacity was being used. Storage in these reservoirs at this time last year was 140 percent of average. First of the month storage in 8 **South Lahontan Region** reservoirs was 299 thousand acre-feet which is 115 percent of average. About 75 percent of available capacity was being used. Storage in these reservoirs at this time last year was 85 percent of average.

RUNOFF Seasonal runoff of streams draining the **North** Lahontan Region totaled 605 thousand acre-feet which is 145 percent of average. Last year, runoff for the same period was 325 percent of average. Seasonal runoff of streams draining the **South Lahontan Region** area totaled 80 thousand acre-feet which is 105 percent of average. Last year, runoff for the same period was 110 percent of average.

SAN FRANCISCO BAY AND CENTRAL COAST REGIONS



PRECIPITATION Seasonal precipitation (October 1 through to the end of April) on the **San Francisco Bay Region** was 70 percent of normal. Precipitation last month was about 160 percent of the monthly average. Season precipitation at this time last year stood at 175 percent of normal. Seasonal precipitation (October 1 through to the end of April) on the **Central Coast Region** was 60 percent of normal. Precipitation last month was about 65 percent of the monthly average. Season precipitation at this time last year stood at 155 percent of normal.

RESERVOIR STORAGE First of the month storage in 17 San Francisco Region 43 reservoirs was 471 thousand acre-feet which is 90 percent of average. About 65 percent of available capacity was being used. Storage in these reservoirs at this time last year was 100 percent of average. First of the month storage in 6 **Central Coast Region** reservoirs was 432 thousand acre-feet which is 60 percent of average. About 45 percent of available capacity was being used. Storage in these reservoirs at this time last year was 100 percent of average.

RUNOFF Seasonal runoff of streams draining the **San** Francisco Region totaled 20 thousand acre-feet million acre-feet which is 30 percent of average. Last year, runoff for the same period was 215 percent of average. Seasonal runoff of streams draining the **Central Coast Region** area totaled 118 thousand acre-feet million acre-feet which is 40 percent of average. Last year, runoff for the same period was 250 percent of average.

SOUTH COAST AND COLORADO RIVER REGIONS

PRECIPITATION - October through April (seasonal) precipitation on the **South Coast Region** was 40 percent of normal. April precipitation was less than 5 percent of the monthly average. Seasonal precipitation at this time last year was 130 percent of normal. Seasonal precipitation on the **Colorado River-Desert Region** was 30 percent of normal. Precipitation during April was 0 percent of average. Seasonal precipitation at this time last year stood at 140 percent of average.

RESERVOIR STORAGE - May 1 storage in 29 major **South Coast Region** reservoirs was 1.2 million acrefeet or 80 percent of average. About 60 percent of available capacity was being used. Storage in these reservoirs at this time last year was 90 percent of average.

RUNOFF - Seasonal runoff from selected **South Coast Region** streams totaled 24 thousand acre-feet which is 15 percent of average. Seasonal runoff from these streams last year was 80 percent of average.

COLORADO RIVER

The April July inflow to Lake Powell is forecast to be 3.0 million acre-feet, which is 42 percent of average. The May 1 snowpack in the Colorado River basin above Lake Powell was 45 percent of average, lowest in the San Juan River headwaters at less than 1 percent and highest in the Upper Green at 70 percent. On May 1 combined storage in Lakes Powell, Mead, Mohave and Havasu was about 25.3 million acre-feet or about 65 percent of average. About 50 percent of available capacity was in use. Last year at this time, these reservoirs were storing 65 percent of average.

MAJOR WATER DISTRIBUTION PROJECTS RESERVOIR STORAGE

(AVERAGES BASED ON 1966-2015 OR PERIOD RECORD)

		AVERAGE_	STORAGE AT END OF April			
RESERVOIR	CAPACITY 1,000 AF	STORAGE 1,000 AF	2017 1,000 AF	2018 1,000 AF	PERCENT AVERAGE	PERCENT CAPACITY
STATE WATER PROJECT	1,000 AF	1,000 AF	1,000 AF	1,000 AF	AVERAGE	CAPACITI
Lake Oroville	3,538	2,857	2,622	2,433	85%	69%
San Luis Reservoir (SWP)	1,062	937	1,032	893	95%	84%
Lake Del Valle	77	39	41	38	96%	49%
Lake Silverwood	78	69	66	68	99%	88%
Pyramid Lake	180	163	166	166	101%	92%
Castaic Lake	325	288	301	283	98%	87%
Perris Lake	131	105	58	95	90%	72%
CENTRAL VALLEY PROJECT						
Trinity Lake	2,448	1,984	2,302	1,930	97%	79%
Lake Shasta	4,552	3,872	4,263	4,195	108%	92%
Whiskeytown Lake	241	233	232	229	98%	95%
Folsom Lake	977	727	724	866	119%	89%
New Melones Reservoir	2,400	1,483	2,002	2,062	139%	86%
Millerton Lake	521	358	261	451	126%	87%
San Luis Reservoir (CVP)	971	839	966	862	103%	89%
COLORADO RIVER PROJECT						
Lake Mead	26,159	18,823	10,420	10,387	55%	40%
Lake Powell	24,322	16,854	12,149	12,669	75%	52%
Lake Mohave	1,810	1,670	1,684	1,682	101%	93%
Lake Havasu	648	587	594	564	96%	87%
EAST BAY MUNICIPAL UTILITY D	DISTRICT					
Pardee Res	204	184	200	205	111%	100%
Camanche Reservoir	417	265	283	383	145%	92%
East Bay (4 res.)	159	134	134	136	101%	86%
CITY AND COUNTY OF SAN FRA	NCISCO					
Hetch-Hetchy Reservoir	360	189	283	325	172%	90%
Cherry Lake	268	176	183	205	116%	76%
Lake Eleanor	29	17	24	23	132%	79%
South Bay/Peninsula (4 res.)	238	173	152	140	81%	59%
CITY OF LOS ANGELES (D.W.P.)						
Lake Crowley	183	124	93	147	119%	80%
Grant Lake	48	26	33	23	86%	48%
Other Aqueduct Storage (6 res.)	238	173	152	140	81%	59%


TELEMETERED SNOW WATER EQUIVALENTS

TELEMETERED SNOW WATER EQUIVALENTS May 1, 2018									
	(A	VERAGES BASED O		CORD) INCHES OF WATER	P EOLIIVAL ENT				
BASIN NAME		APRIL 1		PERCENT	24 HRS	1 WEEK			
STATION NAME	ELEV	AVERAGE	May 1	OF AVERAGE	PREVIOUS	PREVIOUS			
TRINITY RIVER			•						
Shimmy Lake	6400'	40.3	0.0	0.0	0.0	2.5			
Crowder Flat	5100'	-	0.0	-	0.0	0.0			
Highland Lakes	6030' 5650'	29.9 22.4	0.0 0.0	0.0 0.0	0.0 0.0	0.0 0.0			
Mumbo Basin Bonanza King	6450'	40.5	0.0	0.0	-	0.0			
Red Rock Mountain	6700'	39.6	6.4	16.0	7.1	12.0			
Big Flat	5100'	15.8	0.0	0.0	0.0	0.0			
Scott Mountain	5900'	16.0	0.0	0.0	0.0	0.0			
Peterson Flat Middle Boulder 3	7150' 6200'	29.2 28.3	0.0 0.0	0.0 0.0	0.0 0.0	1.3 0.0			
SACRAMENTO RIVER	0200	20.3	0.0	0.0	0.0	0.0			
Blacks Mountain	7050'	12.7	1.6	12.3	1.6	4.0			
Cedar Pass	7100'	18.1	3.8	21.0	3.5	6.9			
Medicine Lake	6700'	32.6	12.6	38.7	12.5	14.8			
Sand Flat	6750'	42.4	-	-	-	-			
Slate Creek Adin Mountain	5700' 6200'	29.0 13.6	0.0 0.7	0.0 5.1	0.0 0.8	0.0 0.0			
Stouts Meadow	5400'	36.0	0.7	0.0	0.0	2.8			
Snow Mountain	5950'	27.0	8.3	30.7	8.4	10.7			
FEATHER RIVER									
Kettle Rock	7300'	25.5	4.7	18.4	4.6	9.5			
Gold Lake	6750'	36.5	18.8	51.6	18.8	20.6			
Bucks Lake	5750'	44.7	6.0	13.4	6.5	11.3			
Harkness Flat Four Trees	6200' 5150'	28.5 20.0	0.0 0.0	0.0 0.0	0.0 0.0	0.0 0.0			
Humbug	6500'	28.0	8.3	29.6	8.4	12.2			
Grizzly Ridge	6900'	29.7	3.8	12.9	4.0	8.5			
Rattlesnake	6100'	14.0	0.0	0.0	0.0	0.0			
Lower Lassen Peak	8250'	-	48.2	-	47.8	49.2			
Pilot Peak	6800'	52.6	1.8	3.4	2.2	6.8			
EEL RIVER Noel Spring	5100'	_	0.0	_	0.0	0.0			
YUBA & AMERICAN RIVERS	3100		0.0		0.0	0.0			
Carson Pass	8353'	-	13.8	-	14.0	18.8			
Lake Lois	8600'	39.5	-	-	-	-			
Forni Ridge	7600'	37.0	2.0	5.3	2.4	8.4			
Silver Lake	7100'	22.7	1.6	7.2	1.6 0.0	4.9			
Blue Canyon Schneiders	5280' 8750'	9.0 34.5	0.0 28.7	0.0 83.1	28.4	0.0 31.8			
Meadow Lake	7200'	55.5	-	-	-	-			
Robbs Powerhouse	5150'	5.2	0.0	0.0	0.0	0.0			
Robinson Cow Camp	6480'	-	3.5	-	3.9	8.5			
Cent Sierra Snow Lab	6900'	33.6	5.1	15.2	5.4	11.1			
Caples Lake Alpha	8000' 7600'	30.9 35.9	6.6 1.2	21.4 3.3	6.7 1.8	10.6 7.9			
Robbs Saddle	5900'	21.4	0.0	0.0	0.0	0.2			
Huysink	6600'	42.6	6.1	14.4	6.2	9.8			
Van Vleck	6700'	35.9	3.9	10.9	4.5	10.0			
Greek Store	5600'	21.0	0.8	4.0	1.1	7.2			
MOKELUMNE & STANISLAUS RIVERS Highland Meadow	8700'	47.9	35.0	73.1	35.1	39.8			
Gianelli Meadow	8400'	55.5	26.9	48.4	26.6	28.6			
Bloods Creek	7200'	35.5	1.2	3.5	1.5	6.4			
Blue Lakes	8000'	33.1	12.9	38.8	13.3	17.1			
Mud Lake	7900'	44.9	-	-	-	-			
Black Springs	6500'	32.0	3.4	10.5	3.7	7.8			
Stanislaus Meadow Deadman Creek	7750' 9250'	47.5 37.2	13.9	29.3	14.4	18.7			
Lower Relief Valley	8100'	41.2	16.0	38.7	16.1	18.9			
TUOLUMNE & MERCED RIVERS									
Dana Meadows	9800'	27.7	10.6	38.4	10.8	14.2			
Horse Meadow	8400'	48.6	34.5	71.0	35.2	40.9			
Tuolumne Meadows	8600'	22.6	1.3	5.8	1.3	3.3			
Slide Canyon Ostrander Lake	9200' 8200'	41.1 34.8	11.9	- 34.3	- 11.9	27.9 15.7			
Gin Flat	7050'	34.6 34.2	-	34.3 -	- 11.9	13.7			
Tenaya Lake	8150'	33.1	6.2	18.8	6.9	15.8			
White Wolf	7900'	-	4.8	-	5.1	10.1			
Lower Kibbie Ridge	6700'	27.4	0.0	0.0	0.0	0.0			
Paradise Meadow	7650'	41.3	11.5	27.8	12.0	17.8			

SAN JOAQUIN RIVER						
Volcanic Knob	10050'	30.1	19.6	65.2	20.0	21.9
Tamarack Summit	7550'	30.5	0.0	0.0	0.0	0.4
Kaiser Point	9200'	37.8	13.0	34.3	13.7	18.8
Huntington Lake	7000'	20.1	0.0	0.0	0.0	8.0
Green Mountain	7900'	30.8	0.0	0.0	0.0	0.0
Poison Ridge	6900'	28.9	0.0	0.0	0.0	0.0
Graveyard Meadow	6900' 9450'	18.8 32.3	0.0 19.0	0.0 58.8	0.0 20.0	0.0 23.5
Agnew Pass Devils Postpile	7569'	32.3	0.0	30.0	0.0	0.0
Chilkoot Meadow	7150'	38.0	0.0	0.0	0.0	3.4
KINGS RIVER	7130	30.0	0.0	0.0	0.0	0.4
Bishop Pass	11200'	34.0	9.7	28.5	9.8	10.1
Blackcap Basin	10300'	34.3	-	-	-	
Mitchell Meadow	9900'	32.9	21.5	65.2	21.4	25.0
Upper Burnt Corral	9700'	34.6	15.7	45.3	15.4	18.8
State Lakes	10300'	29.0	-	-	-	-
West Woodchuck Meadow	9100'	32.8	3.6	11.0	4.0	10.6
Big Meadows	7600'	25.9	-	-	-	-
Charlotte Lake	10400'	27.5	-	-	-	-
KAWEAH & TULE RIVERS						
Farewell Gap	9500'	34.5	-	-	-	-
Giant Forest Quaking Aspen	6650' 7200'	10.0 21.0	0.0 0.0	0.0 0.0	0.0 0.0	0.0 0.0
KERN RIVER	7200	21.0	0.0	0.0	0.0	0.0
Tunnel Guard Station	8900'	15.6	_	_		
Beach Meadows	7650'	11.0	0.0	0.0	0.0	0.0
Upper Tyndall Creek	11400'	27.7	6.4	23.0	6.3	8.6
Casa Vieja Meadows	8300'	20.9	0.0	0.0	0.0	0.0
Pascoes	9150'	24.9	0.0	0.0	0.0	4.0
Wet Meadows	8950'	30.3	0.0	0.0	0.0	0.0
Chagoopa Plateau	10300'	21.8	1.7	7.6	1.3	6.0
Crabtree Meadow	10700'	19.8	-	-	-	-
SURPRISE VALLEY AREA						
Dismal Swamp	7050'	29.2	21.6	74.0	21.0	21.8
TRUCKEE RIVER						
Independence Camp	7000'	21.8	0.0	0.0	0.0	0.5
Independence Lake	8450'	41.4	35.2	85.0	35.0	37.4
Squaw Valley Gold Coast	8200'	46.5	22.8	49.0	23.1	27.1
Truckee 2 Independence Creek	6400' 6500'	14.3 12.7	0.0 0.0	0.0 0.0	0.0 0.0	1.6 0.0
Big Meadows	8700'	25.7	13.5	52.5	13.8	17.1
LAKE TAHOE BASIN	0700	25.1	13.3	32.3	15.0	11.1
Rubicon Peak 2	7500'	29.1	6.2	21.3	6.4	8.7
Tahoe City Cross	6750'	16.0	0.0	0.0	0.0	0.0
Echo Peak 5	7800'	39.5	7.5	19.0	7.7	14.8
Hagans Meadow	8000'	16.5	0.0	0.0	0.0	0.0
Fallen Leaf Lake	6250'	7.0	0.0	0.0	0.0	0.0
Ward Creek 3	6750'	39.4	7.9	20.1	8.1	13.4
Mount Rose Ski Area	8900'	38.5	27.3	70.9	27.0	30.5
Heavenly Valley	8800'	28.1	12.3	43.8	13.3	17.7
Marlette Lake	8000'	21.1	4.5	21.3	5.0	8.9
CARSON RIVER	6150	4.5	0.0	0.0	0.0	0.0
Spratt Creek Horse Meadow	6150' 8400'	4.5 48.6	0.0 34.5	0.0 71.0	0.0 35.2	0.0 40.9
Burnside Lake	8129'	40.0	11.1	71.0	12.2	16.1
Monitor Pass	8350'	-	1.3	-	1.2	8.4
Poison Flat	7900'	16.2	5.2	32.1	5.3	10.3
Forestdale Creek	8017'	-	19.9	-	20.1	23.8
Ebbetts Pass	8700'	38.8	21.6	55.7	22.5	-
WALKER RIVER						
Sonora Pass Bridge	8750'	26.0	9.6	36.9	9.8	12.6
Virginia Lakes Ridge	9300'	20.3	8.6	42.4	9.5	11.5
Lobdell Lake	9200'	17.3	0.0	0.0	1.4	8.0
Summit Meadow	9313'	-	7.9	-	9.2	12.3
Leavitt Leke	7200'	8.0	0.0	0.0	0.0	0.0
Leavitt Lake	9600'	-	45.0	-	44.8	48.8
OWENS RIVER/MONO LAKE Cottonwood Lakes	10150	11 6	0.0	0.0	0.0	0.0
Gem Pass	10150' 10750'	11.6 31.7	0.0	0.0	0.0	0.0
Rock Creek Lakes	9700'	14.0	0.0	0.0	0.0	0.0
South Lake	9600'	16.0	3.0	18.8	3.2	6.4
Big Pine Creek	9800'	17.9	0.0	0.0	0.3	3.8
Sawmill	10200'	19.4	3.1	15.9	3.1	7.5

NORMAL SNOWPACK ACCUMULATION EXPRESSED AS A PERCENT OF APRIL 1ST AVERAGE								
AREA	JANUARY	FEBRUARY	MARCH	APRIL	MAY			
Central Valley North	45%	70%	90%	100%	75%			
Central Valley South	45%	65%	85%	100%	80%			
North Coast	40%	60%	85%	100%	80%			

May 1 Statewide Conditions



SNOWLINES

Next Years Western Snow Conference rotates back to the South Pacific Region. Tentative plans are to meet in the Reno vicinity.

On this months cover is pictured the dramatic effect of the April 8th atmospheric river on the snowpack near New Grace Meadow at an elevation of 8300 feet. The picture was taken on May 3 by Mark Fincher, Wilderness Specialist, Yosemite National Park